

SYSTEMATIC INDEX

(vol. 1, no. 1 - vol. 2, no. 4: 1997-1998)

PALEOZOOLOGY

Deuterostomes

- | | vol. (no.) page |
|--|-----------------|
| Bergström, Jan : Origin of high-rank groups of organisms | 1(1) 1 |

Protoctista

- | | |
|--|----------|
| Hanagata, Satoshi : <i>Haplophragmoides yoshidai</i> , a new name for a homonym of a Japanese foraminiferal species | 2(2) 148 |
| Ling, Hsin Yi and Haddock, Steven, H.D. : The enclosing latticed sphere of <i>Tuscaridium cygneum</i> (Murray), a eurybathyal phaeodarian Radiolaria, from the North Pacific | 1(2) 144 |
| Sugiyama, Kazuhiro and Anderson, O. Roger : The fine structures of some living <i>Spyrida</i> (Nassellaria, Radiolaria) and their implications for nassellarian classification | 2(2) 75 |
| Takemoto, Ayumi and Oda, Motoyoshi : New planktic foraminiferal transfer functions for the Kuroshio-Oyashio Current region off Japan | 1(4) 291 |
| Takemura, Atsushi and Ling, Hsin Yi : Taxonomy and phylogeny of the genus <i>Theocorys</i> (Nassellaria, Radiolaria) from the Eocene and Oligocene sequences in the Antarctic region | 2(3) 155 |
| Umeda, Masaki : Early to Middle Devonian Ceratoikiscidae (Radiolaria) from the Yokokurayama Group in the Kurosegawa Terrane, Southwest Japan | 2(2) 96 |
| Yi, Songsuk, Yun, Hyesu and Yoon, Sun : Calcareous nannoplankton from the Seoguipo Formation of Cheju Island, Korea and its paleoceanographic implications | 2(4) 253 |

Mollusca

- | | |
|---|----------|
| Amano, Kazutaka : Biogeography of the genus <i>Neptunea</i> (Gastropoda : Buccinidae) from the Pliocene and the lower Pleistocene of the Japan Sea borderland | 1(4) 274 |
| Amano, Kazutaka and Vermeij, Geerat J. : Taxonomy and evolution of the genus <i>Ocinebrellus</i> (Gastropoda : Muricidae) in Japan | 2(3) 199 |
| Ehiro, Masayuki and Araki, Hideo : Permian cephalopods of Kurosawa, Kesenuma City in the Southern Kitakami Massif, Northeast Japan | 1(1) 55 |
| Gurung, Damayanti, Takayasu, Katsumi and Matsuoka, Keiji : Middle Miocene-Pliocene freshwater gastropods of the Churia Group, west-central Nepal | 1(3) 166 |
| Honda, Yutaka, Ushiro, Seisuke and Moritani, Sayuri : Miocene mollusks from the Kumano Group of the Ukui area, southeastern part of the Kii Peninsula, southwestern Honshu, Japan | 2(1) 12 |
| Inoue, Keisuke, Ozawa, Tomowo, Nobuhara, Takami and Tomida, Susumu : Reexamination of the Sagara Fauna — Middle Miocene molluscan assemblage from the Sugegaya Formation, Sagara Group, Shizuoka Prefecture, Central Japan — | 1(2) 110 |
| Matsumoto, Tatsuro and Egashira, Naoko : An interesting mid-Cretaceous ammonite from Hokkaido (Studies of the Cretaceous ammonites from Hokkaido and Sakhalin-LXXIX) | 1(2) 150 |
| Matsumoto, Tatsuro, Kawabe, Fumihisa and Kawashita, Yoshitaro : Two ammonite species of <i>Mortonicerias</i> from the Yubari Mountains (Hokkaido) and their geological implications (Studies of the Cretaceous ammonites from Hokkaido and Sakhalin-LXXXII) | 2(3) 170 |
| Matsumoto, Tatsuro and Kawashita, Yoshitaro : Two ammonites species of the genus <i>Sharpeicerias</i> from the Cretaceous of Hokkaido (Studies of Cretaceous ammonites from Hokkaido and Sakhalin-LXXXI) | 2(2) 89 |
| Matsumoto, Tatsuro, Yokoi, Katsujo and Kawashita, Yoshitaro : Further notes on the ammonoid genus <i>Parajaubertella</i> (Studies of Cretaceous Ammonites from Hokkaido and Sakhalin-LXXX) | 1(3) 188 |
| Niko, Shuji : First discosorid cephalopod from Japan | 2(2) 145 |
| Niko, Shuji : Devonian tentaculitids from Central and Northeast Japan | 2(4) 279 |
| Niko, Shuji, Nishida, Tamio and Kyuma, Yuko : Moscovian (Carboniferous) orthoconic cephalopods from Guizhou and Guangxi, South China | 1(2) 100 |
| Niko, Shuji, Pillecuit, Alain and Stampfli, Gerard : Nautiloid and bactritoid cephalopods from the Carboniferous of the | |

Jebel Qamar South area, United Arab Emirates	1(3)	157
Niko, Shuji and Ozawa, Tomowo: Late Gzhelian (Carboniferous) to early Asselian (Permian) non-ammonoid cephalopods from the Taishaku Limestone Group, Southwest Japan	1(1)	47
Okamoto, Takashi and Shibata, Makoto: A cyclic mode of shell growth and its implications in a Late Cretaceous heteromorph ammonite <i>Polyptychoceras pseudogaultinum</i> (Yokoyama)	1(1)	29
Sato, Shin'ichi: Shell microgrowth patterns of bivalves reflecting seasonal change of phytoplankton abundance	1(4)	260
Tanabe, Kazushige, Landman, Neil H. and Mapes, Royal H.: Muscle attachment scars in a Carboniferous goniatite	2(2)	130
Ubukata, Takao: Mantle kinematics and formation of commarginal shell sculpture in <i>Bivalvia</i>	1(2)	132
Yoshida, Katsuhiko: Formation of plications in the Miocene bivalve <i>Mytilus (Plicatomytilus) ksakurai</i> as a consequence of architectural constraint	2(4)	224

Brachiopoda

MacKinnon, David I., Saito, Michiko and Endo, Kazuyoshi: Morphology and systematics of the Recent Japanese brachiopod <i>Shimodaia pterygiota</i> gen. et sp. nov. (Laqueidae: Terebratulida)	1(3)	225
Shen, Shu-Zhong and Tazawa, Jun-ichi: Two permianellids (Brachiopoda) from the Middle Permian of the Southern Kitakami Mountains, Northeast Japan	1(4)	285
Tazawa, Jun-ichi, Ono, Teruo and Hori, Masakazu: Two Permian lytoniid brachiopods from Akasaka, central Japan	2(4)	239

Arthropoda

Chen, Pei-Ji and Suzuki, Shigeyuki: A new species of <i>Nemestheria</i> (Crustacea, Conchostraca) from the Aioi Group (Cretaceous) of Okayama, Japan	2(1)	25
Irizuki, Toshiaki, Ishizaki, Kunihiro, Takahashi, Masaki and Usami, Morihiro: Ostracode faunal changes after the mid-Neogene climatic optimum elucidated in the Middle Miocene Kobana Formation, Central Japan	2(1)	30
Karasawa, Hiroaki: Two new species of Decapoda (Crustacea) from the upper Cretaceous Izumi Group, Japan	2(4)	217

Bryozoa

Nishizawa, Yasuo and Sakagami, Sumio: Phylogenetic significance of a new cheilostome bryozoan species, <i>Dysnoetocella? voigti</i> from the Upper Cretaceous Izumi Group in Shikoku, Japan	1(4)	267
Sakagami, Sumio and Pillevuit, Alain: Permian bryozoans from the exotic formations in Oman	1(3)	200

Echinodermata

Donovan, Stephen K.: Survival of crinoid stalk fragments and its taphonomic implications: discussion	2(4)	283
Ishida, Yoshiaki and Kurita, Yoshitaka: <i>Ophiura sarsii sarsii</i> (Echinodermata, Ophiuroidea) from the Late Pliocene Hachioji Formation in Niigata Prefecture, Central Japan	2(2)	137
Oji, Tatsuo and Amemiya, Shonan: Survival of crinoid stalk fragments and its taphonomic implications	2(1)	67
Oji, Tatsuo and Amemiya, Shonan: Survival of crinoid stalk fragments and its taphonomic implications: additional discussion	2(4)	285

Conodonts

Koike, Toshio: Triassic coniform conodont genera <i>Aduncodina</i> and <i>Neostrachanognathus</i>	2(2)	120
Mizuno, Yoshihiro: Conodont faunas across the Mid-Carboniferous boundary in the Hina Limestone, Southwest Japan	1(4)	237

Vertebrata

Cook, Elizabeth, Isaji, Shinji and Manabe, Makoto: Preliminary results of a taphonomic study of a vertebrate accumulation from the Tetori Group (Lower Cretaceous) of Japan	2(1)	47
Evans, Susan E. and Manabe, Makoto: Early Cretaceous frog remains from the Okurodani Formation, Tetori Group, Japan	2(4)	275
Forsten, Ann: A note on the Jinomori horse, <i>Equus nipponicus</i> Shikama and Onuki (Mammalia, Equidae), and some		

other equine remains from Japan	1(4)	316
Jin, Chang-zhu and Kawamura, Yoshinari: A new species of the extinct shrew <i>Paenelimnoecus</i> from the Pliocene of Yinan, Shandong Province, northern China	1(1)	67
Kohno, Naoki: The first record of an amphicyonid (Mammalia: Carnivora) from Japan, and its implication for amphicyonid paleobiogeography	1(4)	311
Kuwayama, Ryu, Ozawa, Tomowo and Otsuka, Hiroyuki: Fossil red deer <i>Cervus elaphus</i> from the sea-floor of the East China Sea off Amakusa-shimoshima Island, southwestern Kyushu, Japan	2(1)	1
Matsuoka, Hiroshige, Sakakura, Fujio and Ohe, Fumio: A Miocene pseudodontorn (Pelecaniformes: Pelagornithidae) from the Ichishi Group of Misato, Mie Prefecture, Central Japan	2(4)	246
Miyata, Kazunori and Tomida, Yukimitsu: A new tillodont from the early Middle Eocene of Japan and its implication to the subfamily Trogosinae (Tillodontia: Mammalia)	2(1)	53
Miyata, Kazunori and Tomida, Yukimitsu: <i>Trogosus</i> -like tillodont (Tillodontia, Mammalia) from the early Middle Eocene of Japan	2(3)	193

Trace fossil

Kotake, Nobuhiro: Ethological interpretation of the trace fossil <i>Zoophycos</i> in the Hikoroichi Formation (Lower Carboniferous), southern Kitakami Mountains, Northeast Japan	1(1)	15
Miller, William, III and Aalto, Kenneth R.: Anatomy of a complex trace fossil: <i>Phymatoderma</i> from Pliocene bathyal mudstone, northwestern Ecuador	2(4)	266

PALEOBOTANY

Paleobotany

Matsumoto, Midori, Ohsawa, Takeshi, Nishida, Makoto and Nishida, Harufumi: <i>Glyptostrobus rubenosawaensis</i> sp. nov., a new permineralized conifer species from the Middle Miocene, Central Hokkaido, Japan	1(2)	81
Ohana, Tamiko, Kimura, Tatsuaki and Chitaley, Shya: <i>Bennetticarpus yezoites</i> sp. nov. (Bennettitales) from the Upper Cretaceous of Hokkaido, Japan	2(2)	108
Saiki, Ken'ichi: <i>Frenelopsis pombetsuensis</i> : a new cheirolepidiaceous conifer from the Lower Cretaceous (Albian) of Hokkaido, Japan	1(2)	126
Takimoto, Hideo, Ohana, Tamiko and Kimura, Tatsuaki: Two new <i>Nilssoniocladus</i> species from the Jurassic (Oxfordian) Tochikubo Formation, Northeast Honshu, Japan	1(3)	180

Dinophyta

Matsuoka, Kazumi, Pholpunthin, Pornsilp and Fukuyo, Yasuwo: Is the archeopyle of <i>Tuberculodinium vancampoeae</i> (Rossignol) (Gonyaulacales, Dinophyceae) on the hypocyst?	2(3)	183
---	------	-----

INDEX OF GENERA AND SPECIES

Notes : Words listed are names of genus and species-groups which are described, discussed or illustrated in the articles in Paleontological Research, vol.1, no.1 to vol.2, no.4. Words in bold-type are names of newly proposed taxa.

A

- vol. (no.), page, fig(s).
- Acila divaricata submirabilis* 1(2), 114, fig. 5
Aduncodina 2(2), 125
Aduncodina unicastata 2(2), 125, fig. 8
Alternifenestella cf. *A. horologia* 1(3), 214, fig. 5
Alternifenestella sp. A 1(3), 215, fig. 6
Alternifenestella sp. B 1(3), 215, fig. 6
Alternifenestella sp. C 1(3), 215, fig. 6
Alternifenestella sp. D 1(3), 216, fig. 6
Angulyagra 1(3), 171
Angulyagra sp. 1(3), 171, fig. 4
Anisocorbula venusta 1(2), 118, fig. 5

B

- Bactrites* cf. *nagatoensis* 1(2), 106, figs. 3, 4
Bactrites cf. *quadrilineatus* 1(3), 164, fig. 3
Bactrites faqingensis 1(2), 104, fig. 4
Bactrites sp. 1(1), 52, fig. 3
Bactrites sp. 1(2), 106, fig. 4
Bellamya 1(3), 168
Bellamya celsispiralis 1(3), 169, fig. 4
Bellamya sp. A 1(3), 169, fig. 4
Bellamya sp. B 1(3), 169, fig. 4
Bennetticarpus 2(2), 108
Bennetticarpus yezoites 2(2), 108, figs. 1-6
Bhimaites 1(2), 150
Bhimaites ? kawai 1(2), 150, figs. 1-3
Bithynia 1(3), 172
Bithynia sp. 1(3), 172, fig. 5
Bogoslovskya guizhouensis 1(2), 100, fig. 2
Bogoslovskya miharanoroensis 1(1), 48, fig. 2
Brotia 1(3), 173
Brotia palaeocostula 1(3), 173, fig. 5
Brotia sp. A 1(3), 174, fig. 6
Brotia sp. B 1(3), 174, fig. 6
Brotia sp. C 1(3), 176, fig. 6

C

- "*Callianassa*" (s.l.) *masanorii*** 2(4), 220, fig. 3
Callista chinensis 1(2), 118, fig. 5
Calocyclus 2(3), 159
Calocyclus ? nakasekoi 2(3), 160, figs. 3, 5
Capulus ? sp. 1(2), 119, fig. 5
Ceratoikiscum lyratum 2(2), 98, fig. 5
Ceratoikiscum turgidum 2(2), 98, fig. 5
Cervus elaphus 2(1), 1, figs. 3, 4

- Circulaforma* sp. aff. *C. sp. A* 2(2), 103, fig. 7
Conus (Chelyconus) tokunagai 1(2), 121, fig. 5
Conus (Endemoconus) sp. cf. C. (E.) sieboldi 1(2), 121, fig. 5
Coscinophora magnifica 2(4), 241, fig. 2
Crenulilimopsis sp. cf. *C. oblonga* 1(2), 116, fig. 5
Ctenobactrites ? sp. 1(2), 107, fig. 3

D

- Declinognathodus inaequalis* 1(4), 248, fig. 12
Declinognathodus japonicus 1(4), 249, fig. 12
Declinognathodus noduliferus 1(4), 250, fig. 12
Dosinia (Phacosoma) kawagensis 2(1), 20, fig. 5
Dyscritella tenuirama 1(3), 208, fig. 3
Dyscritella sp. 1(3), 210, fig. 4
Dysnoetocella 1(4), 267
Dysnoetocella ? voigti 1(4), 268, figs. 2, 3

E

- Equus nipponicus* 1(4), 316
Eridopora parasitica 1(3), 206, fig. 2
Eumedicottia primas 1(1), 64, figs. 4, 9

F

- Fabifenestella subpermiana* 1(3), 217, fig. 7
Fistulipora cf. *F. wanneri* 1(3), 206, fig. 2
Frenelopsis pombetsuensis 1(2), 127, figs. 2, 3
Fulgoraria (Musashia) yanagidaniensis 2(1), 21, fig. 5

G

- Geisonocerina ? sp.* 1(1), 50, fig. 2
Girtpora cf. *G. ramosa* 1(3), 214, fig. 5
Glanta 2(2), 98
Glanta fragilis 2(2), 99, fig. 5
Glanta yokokurayamaensis 2(2), 101, fig. 5
Glanta sp. 2(2), 101, fig. 5
Glycymeris idensis 1(2), 116, fig. 5
Glycymeris izumoensis 1(2), 116, fig. 5
Glyptostrobus rubenosawaensis 1(2), 87, figs. 5-11
Gnathodus bilineatus 1(4), 250, fig. 11
Goniocladia timorensis 1(3), 208, fig. 3
Gotlandellites fukujiensis 2(4), 281, fig. 1
Gyraulus 1(3), 178
Gyraulus sp. 1(3), 178, fig. 6

H

- Haplophragmoides kushiroensis* 2(2), 148
Haplophragmoides yoshidai 2(2), 148
Helenifore sp. A 2(2), 104, fig. 8
Hexagonella kobayashii 1(3), 206, figs. 2, 3
Higotherium 2(1), 55
Higotherium hypsodon 2(1), 55, figs. 2–4
Hoploparia miyamotoi 2(4), 217, figs. 1, 2

I

- Indoplanorbis* 1(3), 176
Indoplanorbis cf. *exustus* 1(3), 176, fig. 6

J

- Jiingites kesennumensis** 1(1), 60, figs. 4, 5

L

- Laterispina* 1(4), 289
Laterispina parallela 1(4), 289, fig. 5
Laxifenestella lahuseni 1(3), 216, fig. 6
Leptodus nobilis 2(4), 241, figs. 2, 3
Lochriea commutata 1(4), 251, fig. 11
Lochriea nodosa 1(4), 253, fig. 11
Lopingoceras hayasakai 1(1), 50, fig. 3
Lymnaea 1(3), 176
Lymnaea sp. 1(3), 176, fig. 6

M

- Megacardita ferruginosa* 1(2), 117, fig. 5
Melanoides 1(3), 172
Melanoides cf. *tuberculata* 1(3), 173, fig. 5
Mericoceras guangxiense 1(2), 101, fig. 3
Mericoceras guangxiense? 1(2), 104, fig. 3
Mericoceras sp. 1(2), 104, fig. 3
Michelinoceras sp. 1 1(3), 158, fig. 3
Michelinoceras ? sp. 1(1), 48, fig. 2
Michelinoceras ? sp. 2 1(3), 159, fig. 3
Mimogeisonoceras ? sp. 1(2), 104, fig. 4
Minilya duplaris 1(3), 216, fig. 6
Minilya sp. 1(3), 216, fig. 6
Mitrothoceras ? sp. 1(3), 161, fig. 5
Mooreoceras ? sp. 1 1(3), 161, fig. 4
Mooreoceras ? sp. 2 1(3), 161, fig. 5
Mortoniceras (*Mortoniceras*) 2(3), 172
Mortoniceras (*Mortoniceras*) cf. *geometricum* 2(3), 172, fig. 3
Mortoniceras (*Mortoniceras*) *rostratum* 2(3), 176, figs. 4–7
Mytilus (*Plicatomytilus*) *ksakurai* 2(4), 225, fig. 1

N

- Nemestheria* 2(1), 25
Nemestheria aidaensis 2(1), 28, figs. 3, 4
Neognathodus symmetricus 1(4), 253, fig. 12
Neolochriea 1(4), 253

- Neolochriea hisaharui*** 1(4), 255, fig. 13
Neolochriea hisayoshii 1(4), 255, fig. 13
Neolochriea koikei 1(4), 256, fig. 13
Neolochriea nagatoensis 1(4), 256, fig. 13
Neostrachanognathus 2(2), 125
Neostrachanognathus tahoensis 2(2), 127, fig. 9
Neptunea (*Golikovia*) 1(4), 276
Neptunea (*Golikovia*) *nikkoensis* 1(4), 276, fig. 2
Neptunea (*Neptunea*) 1(4), 276
Neptunea (*Neptunea*) *eos* 1(4), 276, fig. 2
Neptunea (*Neptunea*) *hataii* 1(4), 276, figs. 2, 3
Neverita coticae 1(2), 119, fig. 5
Nilssoniocladus 1(3), 182
Nilssoniocladus japonicus 1(3), 183, figs. 1, 3, 4
Nilssoniocladus tairae 1(3), 182, figs. 1–3
Nipponocrassatella nana 1(2), 117, fig. 5
Nipponolimopsis sp. cf. *N. azumana* 1(2), 116, fig. 5

O

- “*Ocenebra*” *katayamai* 2(3), 207, fig. 3
Ocenebrellus 2(3), 200
Ocenebrellus aduncus 2(3), 206, fig. 4
Ocenebrellus inornatus 2(3), 201, fig. 3
Ocenebrellus lumarius 2(3), 204, fig. 3
Ocenebrellus ogasawarai 2(3), 204, figs. 3, 4
Ocenebrellus protoaduncus 2(3), 204, fig. 3
Olivella fulgurata 1(2), 120, fig. 5
Ophiura sarsii sarsii 2(2), 138, figs. 2, 3
Osteodontornis sp. 2(4), 246, figs. 2, 3
Otukaia sp. 1(2), 119, fig. 5

P

- Paenelimnoecus* 1(1), 67
Paenelimnoecus chinensis 1(1), 67, figs. 1–6
Parachouteauoceras 1(1), 52
Parachouteauoceras bingoense 1(1), 52, fig. 3
Parachouteauoceras ? sp. 1(1), 52, fig. 3
Parajaubertella 1(3), 189
Parajaubertella kawakitana 1(3), 189, figs. 3–6
Parajaubertella zizoh 1(3), 194, figs. 7–9
Paralioclema eplicatum 1(3), 210, fig. 4
Penniretepora sp. 1(3), 218, fig. 8
Periploma (*Aelga*) *mitsuganoense* 2(1), 20, fig. 5
Permianella 1(4), 285
Permianella typica 1(4), 288, figs. 2–5
Phalium yokoyamai 1(2), 119, fig. 5
Phos miyagiensis 1(2), 120, fig. 5
Phymatoderma granulatum 2(4), 267, fig. 1
Pila 1(3), 171
Pila sp. 1(3), 171, fig. 5
Pleuonautilus sp. 1(1), 59, figs. 2, 3
Polypora cf. *P. elliptica* 1(3), 220, figs. 6, 8
Polypora cf. *P. gigantea* 1(3), 220, fig. 8
Polypora sp. 1(3), 220, fig. 7
Poterioceratidae gen. et sp. indet. 1(3), 164, fig. 3
Propinacoceras sp. 1(1), 64, fig. 9
Protoholoeciscus 2(2), 103

Protoholoeciscus hindea 2(2), 103, fig. 7
Protoholoeciscus ochiensis 2(2), 104, fig. 8
Protoholoeciscus spinosus 2(2), 104, fig. 7
Protoholoeciscus triangularis 2(2), 104, fig. 8
Protoholoeciscus sp. 2(2), 103, fig. 7
Protoretepora ? sp. 1(3), 222, fig. 8
Pseudagathiceras 1(1), 62
Pseudagathiceras ornatum 1(1), 62, figs. 4, 5

R

Rhombopora aff. *R. lepidodendroides* 1(3), 210, fig. 4
Rhombopora sp. 1(3), 212, fig. 4
Rugofenestella basleoensis 1(3), 217, fig. 7
Rugofenestella omanica 1(3), 218, fig. 7

S

Septopora sp. 1(3), 218, fig. 8
Sharpeiceras 2(2), 89
Sharpeiceras kikuae 2(2), 92, figs. 3, 4
Sharpeiceras mexicanum 2(2), 89, figs. 1, 2
Shimodaia 1(3), 226
Shimodaia pterygiota 1(3), 229, figs. 2, 3, 7
Sinobactrites 1(2), 107
Sinobactrites wuae 1(2), 107, figs. 5, 6
Siphonalia sp. 1(2), 120, fig. 5
Solariella (Machaeroplax) sp. 1(2), 119, fig. 5
Solariella sp. 1(2), 119, fig. 5
Spyroceratinae gen. et sp. indet. 1(3), 163, fig. 5
Stacheoceras sp. 1(1), 61, figs. 4, 5
Stearoceras sp. 1(1), 59, figs. 2, 3

Streblascopora delicatula 1(3), 212, figs. 4, 5
Streblascopora exillis 1(3), 212, fig. 5

T

Tainoceras carinatum 1(1), 57, fig. 3
Tainoceras sp. 1(1), 59, figs. 2, 3
Temperoceras ayimense 1(3), 159, fig. 4
Tentaculites sp. 2(4), 279, fig. 1
Theocorys 2(3), 160
Theocorys kerguelensis 2(3), 160, figs. 3, 5
Theocorys minuta 2(3), 162, figs. 3, 5
Theocorys robusta 2(3), 164, figs. 4, 5
Theocorys saginata 2(3), 164, figs. 4, 5
Theocorys semipolita 2(3), 166, figs. 4, 5
Thracia watanabei 2(1), 20, fig. 5
Timanodictya cf. *T. dichotoma* 1(3), 214, fig. 5
Timorites takaizumii 1(1), 62, figs. 6, 8
Tractabilis ? sp. 2(4), 281, fig. 1

Cf. *Trogosus* sp. 2(3), 194, figs. 2, 3
Turritella (Hataiella) sagai 2(1), 21, fig. 5
Tuscaridium 1(2), 146
Tuscaridium cygneum 1(2), 146, figs. 1, 2

U

Ukhtoceras hidense 2(2), 145, fig. 1

Y

Ysengrinia sp. 1(4), 311, fig. 1

CONTENTS

Vol. 1 No. 1 (published April 30, 1997)

	Page	Figures
Jan Bergström : Origin of high-rank groups of organisms	1- 14	1- 7
Nobuhiro Kotake : Ethological interpretation of the trace fossil <i>Zoophycos</i> in the Hikoroichi Formation (Lower Carboniferous), southern Kitakami Mountains, Northeast Japan	15- 28	1-13
Takashi Okamoto and Makoto Shibata : A cyclic mode of shell growth and its implications in a Late Cretaceous heteromorph ammonite <i>Polyptychoceras pseudogaultinum</i> (Yokoyama)	29- 46	1-15
Shuji Niko and Tomowo Ozawa : Late Gzhelian (Carboniferous) to early Asselian (Permian) non-ammonoid cephalopods from the Taishaku Limestone Group, Southwest Japan	47- 54	1- 3
Masayuki Ehiro and Hideo Araki : Permian cephalopods of Kurosawa, Kesenuma City in the Southern Kitakami Massif, Northeast Japan	55- 66	1- 9
Chang-zhu Jin and Yoshinari Kawamura : A new species of the extinct shrew <i>Paenelimnoecus</i> from the Pliocene of Yinan, Shandong Province, northern China	67- 75	1- 6
PROCEEDINGS	76- 79	

Vol. 1 No. 2 (published June 30, 1997)

Midori Matsumoto, Takeshi Asakawa Ohsawa, Makoto Nishida and Harufumi Nishida : <i>Glyptostrobus rubenosawaensis</i> sp. nov., a new permineralized conifer species from the Middle Miocene, Central Hokkaido, Japan	81- 99	1-12
Shuji Niko, Tamio Nishida and Yuko Kyuma : Moscovian (Carboniferous) orthoconic cephalopods from Guizhou and Guangxi, South China	100-109	1- 6
Keisuke Inoue, Tomowo Ozawa, Takami Nobuhara and Susumu Tomida : Reexamination of the Sagara Fauna — Middle Miocene molluscan assemblage from the Sugegaya Formation, Sagara Group, Shizuoka Prefecture, Central Japan —	110-125	1- 5
Ken'ichi Saiki : <i>Frenelopsis pombetsuensis</i> : a new cheirolepidiaceus conifer from the Lower Cretaceous (Albian) of Hokkaido, Japan	126-131	1- 3
Takao Ubukata : Mantle kinematics and formation of commarginal shell sculpture in <i>Bivalvia</i>	132-143	1- 7
Hsin Yi Ling and Steven H.D. Haddock : The enclosing latticed sphere of <i>Tuscardium cygneum</i> (Murray), a eurybathyal phaeodarian Radiolaria, from the North Pacific	144-149	1- 2

SHORT NOTES

Tatsuro Matsumoto and Naoko Egashira : An interesting mid-Cretaceous ammonite from Hokkaido (Studies of the Cretaceous ammonites from Hokkaido and Sakhalin-LXXIX)	150-152	1- 3
PROCEEDING	153-155	

Vol. 1 No. 3 (published September 30, 1997)

Shuji Niko, Alain Pillevuit and Gerard Stampfli : Nautiloid and bactritoid cephalopods from the Carboniferous of the Jebel Qamar South area, United Arab Emirates	157-165	1- 5
Damayanti Gurung, Katsumi Takayasu and Keiji Matsuoka : Middle Miocene-Pliocene freshwater gastropods of the Churia Group, west-central Nepal	166-179	1- 6
Hideo Takimoto, Tamiko Ohana and Tatsuaki Kimura : Two new <i>Nilssoniocladus</i> species from the Jurassic (Oxfordian) Tochikubo Formation, Northeast Honshu, Japan	180-187	1- 4
Tatsuro Matsumoto, Katsujo Yokoi and Yoshitaro Kawashita : Further notes on the ammonoid genus <i>Parajaubertella</i> (Studies of Cretaceous Ammonites from Hokkaido and Sakhalin-LXXX)	188-199	1- 9
Sumio Sakagami and Alain Pillevuit : Permian bryozoans from the exotic formations in Oman	200-224	1- 8

David I. MacKinnon, Michiko Saito and Kazuyoshi Endo: Morphology and systematics of the Recent Japanese brachiopod <i>Shimodaia pterygiota</i> gen. et sp. nov. (Laqueidae: Terebratulida)	225-233	1- 7
PROCEEDINGS	234-236	

Vol. 1 No.4 (published December 30, 1997)

Yoshihiro Mizuno: Conodont faunas across the Mid-Carboniferous boundary in Hina Limestone, Southwest Japan.	237-259	1-13
Shin'ichi Sato: Shell microgrowth patterns of bivalves reflecting seasonal change of phytoplankton abundance.	260-266	1- 6
Yasuo Nishizawa and Sumio Sakagami: Phylogenetic significance of a new cheilostome bryozoan species, <i>Dysnoetocella? voigti</i> from the Upper Cretaceous Izumi Group in Shikoku, Japan	267-273	1- 3
Kazutaka Amano: Biogeography of the genus <i>Neptunea</i> (Gastropoda: Buccinidae) from the Pliocene and the lower Pleistocene of the Japan Sea borderland.	274-284	1- 4
Shu-Zhong Shen and Jun-ichi Tazawa: Two permianellids (Brachiopoda) from the Middle Permian of the Southern Kitakami Mountains, Northeast Japan.	285-290	1- 5
Ayumi Takemoto and Motoyoshi Oda: New planktic foraminiferal transfer functions for the Kuroshio-Oyashio Current region off Japan	291-310	1-10
Naoki Kohno: The first record of an amphicyonid (Mammalia: Carnivora) from Japan, and its implication for amphicyonid paleobiogeography	311-315	1- 2

SHORT NOTES

Ann Forsten: A note on the Jinomori horse, <i>Equus nipponicus</i> Shikama and Onuki (Mammalia, Equidae), and some other equine remains from Japan	316-317	
--	---------	--

PROCEEDINGS	318-320	
------------------------------	---------	--

Vol. 2 No. 1 (published April 30, 1998)

Ryu Kuwayama, Tomowo Ozawa and Hiroyuki Otsuka: Fossil red deer <i>Cervus elaphus</i> from the sea floor of the East China Sea off Amakusa-shimoshima Island, southwestern Kyushu, Japan	1- 11	1- 8
Yutaka Honda, Seisuke Ushiro and Sayuri Moritani: Miocene mollusks from the Kumano Group of the Ukui area, southeastern part of the Kii Peninsula, southwestern Honshu, Japan	12- 24	1- 5
Pei-Ji Chen and Shigeyuki Suzuki: A new species of <i>Nemestheria</i> (Crustacea, Conchostraca) from the Aioi Group (Cretaceous) of Okayama, Japan	25- 29	1- 4
Toshiaki Irizuki, Kunihiro Ishizaki, Masaki Takahashi and Morihiro Usami: Ostracode faunal changes after the mid-Neogene climatic optimum elucidated in the Middle Miocene Kobana Formation, Central Japan	30- 46	1- 9
Elizabeth Cook, Shinji Isaji and Makoto Manabe: Preliminary results of a taphonomic study of a vertebrate accumulation from the Tetori Group (Lower Cretaceous) of Japan.	47- 52	1- 4
Kazunori Miyata and Yukimitsu Tomida: A new tillodont from the early Middle Eocene of Japan and its implication to the subfamily Trogosinae (Tillodontia: Mammalia)	53- 66	1- 7

SHORT NOTES

Tatsuo Oji and Shonan Amemiya: Survival of crinoid stalk fragments and its taphonomic implications	67- 70	1- 2
--	--------	------

PROCEEDINGS	71- 73	
------------------------------	--------	--

Vol. 2 No. 2 (published June 30, 1998)

Kazuhiro Sugiyama and O. Roger Anderson: The fine structures of some living Spyrida (Nassellaria, Radiolaria) and their implications for nassellarian classification.	75- 88	1-10
Tatsuro Matsumoto and Yoshitaro Kawashita: Two ammonites of the genus <i>Sharpeiceras</i> from the Cretaceous of Hokkaido (Studies of Cretaceous ammonites from Hokkaido and Sakhalin-LXXXI)	89- 95	1- 4

Masaki Umeda : Early to Middle Devonian Ceratoliskidae (Radiolaria) from the Yokokurayama Group in the Kurosegawa Terrane, Southwest Japan	96-107	1- 9
Tamiko Ohana, Tatsuaki Kimura and Shya Chitaley : <i>Bennetticarpus yezoites</i> sp. nov. (Bennettitales) from the Upper Cretaceous of Hokkaido, Japan	108-119	1- 6
Toshio Koike : Triassic coniform conodont genera <i>Aduncodina</i> and <i>Neostrachanognathus</i>	120-129	1- 9
Kazushige Tanabe, Neil H. Landman and Royal H. Mapes : Muscle attachment scars in a Carboniferous goniatite	130-136	1- 4
Yoshiaki Ishida and Yoshitaka Kurita : <i>Ophiura sarsii sarsii</i> (Echinodermata, Ophiuroidea) from the Late Pliocene Hachioji Formation in Niigata Prefecture, Central Japan	137-144	1- 7

SHORT NOTES

Shuji Niko : First discosorid cephalopod from Japan	145-147	1
Satoshi Hanagata : <i>Haplophragmoides yoshidai</i> , a new name for a homonym of a Japanese foraminiferal species	148-150	

PROCEEDINGS	151-153	
--------------------------	---------	--

Vol. 2 No. 3 (published September 30, 1998)

Atsushi Takemura and Hsin Yi Ling : Taxonomy and phylogeny of the genus <i>Theocorys</i> (Nassellaria, Radiolaria) from the Eocene and Oligocene sequences in the Antarctic region	155-169	1- 6
Tatsuro Matsumoto, Fumihisa Kawabe and Yoshitaro Kawashita : Two ammonite species of <i>Mortonicerus</i> from the Yubari Mountains (Hokkaido) and their geological implications (Studies of the Cretaceous ammonites from Hokkaido and Sakhalin-LXXXII)	170-182	1- 7
Kazumi Matsuoka, Pornsilp Pholpunthin and Yasuwo Fukuyo : Is the archeopyle of <i>Tuberculodinium van-campoae</i> (Rossignol) (Gonyaulacales, Dinophyceae) on the hypocyst ?	183-192	1- 8
Kazunori Miyata and Yukimitsu Tomida : <i>Trogosus</i> -like tillodont (Tillodontia, Mammalia) from the early Middle Eocene of Japan	193-198	1- 3
Kazutaka Amano and Geerat J. Vermei : Taxonomy and evolution of the genus <i>Ocinebrellus</i> (Gastropoda : Muricidae) in Japan	199-212	1- 5
PROCEEDINGS	213-215	

Vol. 2 No. 4 (published December 30, 1998)

Hiroaki Karasawa : Two new species of Decapoda (Crustacea) from the upper Cretaceous Izumi Group, Japan	217-223	1- 3
Katsuhiko Yoshida : Formation of plications in the Miocene bivalve <i>Mytilus (Plicatomytilus) ksakurai</i> as a consequence of architectural constraint	224-238	1-11
Jun-ichi Tazawa, Teruo Ono and Masakazu Hori : Two Permian lyttoniid brachiopods from Akasaka, central Japan	239-245	1- 4
Hiroshige Matsuoka, Fujio Sakakura and Fumio Ohe : A Miocene pseudodontorn (Pelecaniformes : Pelagornithidae) from the Ichishi Group of Misato, Mie Prefecture, Central Japan	246-252	1- 5
Songsuk Yi, Hyesu Yun and Sun Yoon : Calcareous nannoplankton from the Seoguiipo Formation of Cheju Island, Korea and its paleoceanographic implications	253-265	1- 5
William Miller, III and Kenneth R. Aalto : Anatomy of a complex trace fossil : <i>Phymatoderma</i> from Pliocene bathyal mudstone, northwestern Ecuador	266-274	1- 7

SHORT NOTES

Susan E. Evans and Makoto Manabe : Early Cretaceous frog remains from the Okurodani Formation, Tetori Group, Japan	275-278	1- 4
Shuji Niko : Devonian tentaculitids from Central and Northeast Japan	279-282	1
Stephen K. Donovan : Survival of crinoid stalk fragments and its taphonomic implications: discussion	283-284	
Tatsuo Oji and Shonan Amemiya : Survival of crinoid stalk fragments and its taphonomic implications: additional discussion	285-286	

PROCEEDINGS	287-289
SYSTEMATIC INDEX	290-292
INDEX OF GENERA AND SPECIES	293-295

行事予定

- ◎1999年年会は1999年1月29日(金)～1月31日(日)に、「東北大学」で開催されます。シンポジウムとして1月29日に「復元の科学3, 生物事変, 世話人: 海保邦夫, 西 広嗣, 大野照文」が行われます。
- ◎第148回例会は, 1999年6月26日(土)～6月27日(日)に, 「兵庫県立人と自然の博物館」で行われます。一般講演の申し込み締切は5月7日です。シンポジウムの企画をお持ちの方は, 1998年12月末までに行事係までお申し込み下さい。
- ◎2000年年会・総会には現在の所「早稲田大学」から開催の申し込みがありました(決定ではありません)。シンポジウム企画の申し込み締切は1999年3月末日です。
- ◎第149回例会(開催予定時期: 2000年の6月末頃)は, 今の所開催申し込みがありません。開催を計画されている機関がありましたら, お申し込み下さい。
- ◎現在, 常務委員会は2001年からの学会行事の変更を検討しております。2001年からは, 従来とは異なった時期や開催形式で年会, 例会, 総会が開催される可能性があります。従いまして, 第149回例会(2000年6月末に開催予定)までは, 従来通り開催の申し込みを受け付けますが, 2001年以降の開催申し込みは, しばらくの間見あわせて頂きます。

講演の申し込み先: 〒240-8501 横浜市保土ヶ谷区常盤台 79-2

横浜国立大学教育人間科学部自然環境講座 間嶋隆一 (行事係)

問い合わせ先 間嶋隆一: TEL 045-339-3349 直通 FAX 045-339-3264 学部事務室

E-mail majima@ed.ynu.ac.jp HDA01631@niftyserve.or.jp

樽 創 (行事係幹事):

〒250-0031 小田原市入生田 499

神奈川県立生命の星・地球博物館

TEL 0465-21-1515 FAX 0465-23-8846

E-mail taru@pat-net.or.jp

本誌の発行に要する費用は, 会員の会費以外に, 文部省科学研究費補助金ならびに賛助会員からの会費が当てられています。現在の賛助会員は下記の通りです。

インドネシア石油株式会社	神奈川県立生命の星・地球博物館	北九州市立自然史博物館
石油資源開発株式会社	ダイヤコンサルタント	帝国石油株式会社
兵庫県立人と自然の博物館	ミュージアムパーク茨城県自然博物館	(アイウエオ順)

○文部省科学研究費補助金(研究成果公開促進費)による。

<p>1998年12月27日 印刷</p> <p>1998年12月30日 発行</p> <p>ISSN 1342-8144</p> <p>Paleontological Research</p> <p>第2巻, 第4号</p> <p>2,500円</p>	<p style="text-align: center;">発行者</p> <p>日本古生物学会 〒113-8622 東京都文京区本駒込5-16-9 日本学会事務センター内 電話 03-5814-5801</p> <p style="text-align: center;">編集者</p> <p>森 啓・前田晴良</p> <p style="text-align: center;">編集幹事</p> <p>島本昌憲</p> <p style="text-align: center;">印刷者</p> <p>〒984-0011 仙台市若林区六丁の目西町8-45 笹氣出版印刷株式会社 笹氣 幸緒 本社 022-288-5555 東京 03-3455-4415</p>
---	--

Paleontological Research

Vol. 2 No. 4


December 30, 1998

CONTENTS

Hiroaki Karasawa : Two new species of Decapoda (Crustacea) from the Upper Cretaceous Izumi Group, Japan	217
Katsuhiko Yoshida : Formation of plications in the Miocene bivalve <i>Mytilus (Plicatomytilus) ksakurai</i> as a consequence of architectural constraint	224
Jun-ichi Tazawa, Teruo Ono and Masakazu Hori : Two Permian lytoniid brachiopods from Akasaka, central Japan	239
Hiroshige Matsuoka, Fujio Sakakura and Fumio Ohe : A Miocene pseudodontorn (Pelecaniformes: Pelagornithidae) from the Ichishi Group of Misato, Mie Prefecture, Central Japan	246
Songsuk Yi, Hyesu Yun and Sun Yoon : Calcareous nannoplankton from the Seoguipo Formation of Cheju Island, Korea and its paleoceanographic implications	253
William Miller, III and Kenneth R. Aalto : Anatomy of a complex trace fossil: <i>Phymatoderma</i> from Pliocene bathyal mudstone, northwestern Ecuador	266
SHORT NOTES	
Susan E. Evans and Makoto Manabe : Early Cretaceous frog remains from the Okurodani Formation, Tetori Group, Japan	275
Shuji Niko : Devonian tentaculitids from Central and Northeast Japan	279
Stephen K. Donovan : Survival of crinoid stalk fragments and its taphonomic implications: discussion	283
Tatsuo Oji and Shonan Amemiya : Survival of crinoid stalk fragments and its taphonomic implications: additional discussion	285
PROCEEDINGS	287
SYSTEMATIC INDEX	291
INDEX OF GENERA AND SPECIES	294

HECKMAN
B I N D E R Y, I N C.
Bound-To-Pleas®
JUNE 00
N. MANCHESTER, INDIANA 46962

SMITHSONIAN INSTITUTION LIBRARIES



3 9088 01429 0100